

Historical Species List for Booderee Botanic Gardens from Birdata

Common Name	Scientific Name	Count	Reporting Rate
Australasian Grebe	<i>Tachybaptus novaehollandiae</i>	34	27.20%
Australian King-Parrot	<i>Alisterus scapularis</i>	21	16.80%
Australian Magpie	<i>Gymnorhina tibicen</i>	14	11.20%
Australian Raven	<i>Corvus coronoides</i>	28	22.40%
Australian Wood Duck	<i>Chenonetta jubata</i>	10	8.00%
Azure Kingfisher	<i>Ceyx azureus</i>	2	1.60%
Bar-shouldered Dove	<i>Geopelia humeralis</i>	1	0.80%
Bassian Thrush	<i>Zoothera lunulata</i>	29	23.20%
Beautiful Firetail	<i>Stagonopleura bella</i>	1	0.80%
Black Swan	<i>Cygnus atratus</i>	6	4.80%
Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>	14	11.20%
Black-faced Monarch	<i>Monarcha melanopsis</i>	17	13.60%
Black-fronted Dotterel	<i>Elsyornis melanops</i>	1	0.80%
Brown Cuckoo-Dove	<i>Macropygia phasianella</i>	1	0.80%
Brown Falcon	<i>Falco berigora</i>	1	0.80%
Brown Gerygone	<i>Gerygone mouki</i>	56	44.80%
Brown Goshawk	<i>Accipiter fasciatus</i>	1	0.80%
Brown Thornbill	<i>Acanthiza pusilla</i>	72	57.60%
Brown-headed Honeyeater	<i>Melithreptus brevirostris</i>	5	4.00%
Brush Bronzewing	<i>Phaps elegans</i>	1	0.80%
Brush Cuckoo	<i>Cacomantis variolosus</i>	3	2.40%
Buff-rumped Thornbill	<i>Acanthiza reguloides</i>	2	1.60%
Channel-billed Cuckoo	<i>Scythrops novaehollandiae</i>	1	0.80%
Chestnut Teal	<i>Anas castanea</i>	11	8.80%
Chestnut-rumped Heathwren	<i>Calamanthus pyrrhopygius</i>	1	0.80%
Cicadabird	<i>Edolisoma tenuirostris</i>	1	0.80%
Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>	2	1.60%
Common Blackbird	<i>Turdus merula</i>	3	2.40%
Common Bronzewing	<i>Phaps chalcoptera</i>	3	2.40%
Crested Shrike-tit	<i>Falcunculus frontatus</i>	4	3.20%
Crimson Rosella	<i>Platycercus elegans</i>	45	36.00%
Crow & Raven spp		2	1.60%
Dollarbird	<i>Eurystomus orientalis</i>	1	0.80%
Eastern Koel	<i>Eudynamys orientalis</i>	2	1.60%
Eastern Rosella	<i>Platycercus eximius</i>	1	0.80%
Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>	90	72.00%
Eastern Whipbird	<i>Psophodes olivaceus</i>	29	23.20%
Eastern Yellow Robin	<i>Eopsaltria australis</i>	90	72.00%
Eurasian Coot	<i>Fulica atra</i>	1	0.80%
Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>	13	10.40%
Fuscous Honeyeater	<i>Ptilotula fusca</i>	5	4.00%
Galah	<i>Eolophus roseicapilla</i>	3	2.40%
Gang-gang Cockatoo	<i>Callocephalon fimbriatum</i>	14	11.20%
Glossy Black-Cockatoo	<i>Calyptorhynchus lathami</i>	6	4.80%
Golden Whistler	<i>Pachycephala pectoralis</i>	40	32.00%
Great Cormorant	<i>Phalacrocorax carbo</i>	8	6.40%
Grey Butcherbird	<i>Cracticus torquatus</i>	6	4.80%
Grey Fantail	<i>Rhipidura fuliginosa</i>	67	53.60%
Grey Goshawk	<i>Accipiter novaehollandiae</i>	2	1.60%
Grey Shrike-thrush	<i>Colluricincla harmonica</i>	8	6.40%
Grey Teal	<i>Anas gracilis</i>	6	4.80%
Hardhead	<i>Aythya australis</i>	28	22.40%
Hoary-headed Grebe	<i>Poliiocephalus poliocephalus</i>	19	15.20%
Horsfield's Bronze-Cuckoo	<i>Chalcites basalis</i>	1	0.80%
Jacky Winter	<i>Microeca fascinans</i>	1	0.80%
Laughing Kookaburra	<i>Dacelo novaeguineae</i>	67	53.60%
Leaden Flycatcher	<i>Myiagra rubecula</i>	7	5.60%

Lewin's Honeyeater	Meliphaga lewinii	76	60.80%
Little Black Cormorant	Phalacrocorax sulcirostris	1	0.80%
Little Eagle	Hieraaetus morphnoides	2	1.60%
Little Lorikeet	Glossopsitta pusilla	5	4.00%
Little Pied Cormorant	Microcarbo melanoleucos	9	7.20%
Little Wattlebird	Anthochaera chrysoptera	57	45.60%
Masked Lapwing	Vanellus miles	18	14.40%
Mistletoebird	Dicaeum hirundinaceum	1	0.80%
Musk Lorikeet	Glossopsitta concinna	4	3.20%
New Holland Honeyeater	Phylidonyris novaehollandiae	64	51.20%
Noisy Friarbird	Philemon corniculatus	38	30.40%
Noisy Miner	Manorina melanocephala	3	2.40%
Olive-backed Oriole	Oriolus sagittatus	13	10.40%
Pacific Black Duck	Anas superciliosa	30	24.00%
Pied Cormorant	Phalacrocorax varius	1	0.80%
Pied Currawong	Strepera graculina	35	28.00%
Rainbow Lorikeet	Trichoglossus moluccanus	64	51.20%
Red Wattlebird	Anthochaera carunculata	62	49.60%
Red-browed Finch	Neochmia temporalis	38	30.40%
Rose Robin	Petroica rosea	4	3.20%
Rufous Fantail	Rhipidura rufifrons	13	10.40%
Rufous Whistler	Pachycephala rufiventris	13	10.40%
Sacred Kingfisher	Todiramphus sanctus	7	5.60%
Satin Bowerbird	Ptilonorhynchus violaceus	64	51.20%
Satin Flycatcher	Myiagra cyanoleuca	2	1.60%
Scarlet Honeyeater	Myzomela sanguinolenta	25	20.00%
Scarlet Robin	Petroica multicolor	1	0.80%
Shining Bronze-Cuckoo	Chalcites lucidus	3	2.40%
Silvereye	Zosterops lateralis	47	37.60%
Southern Boobook	Ninox boobook	1	0.80%
Southern Emu-wren	Stipiturus malachurus	1	0.80%
Spotted Pardalote	Pardalotus punctatus	56	44.80%
Square-tailed Kite	Lophoictinia isura	1	0.80%
Striated Pardalote	Pardalotus striatus	3	2.40%
Striated Thornbill	Acanthiza lineata	19	15.20%
Sulphur-crested Cockatoo	Cacatua galerita	1	0.80%
Superb Fairy-wren	Malurus cyaneus	65	52.00%
Swamp Harrier	Circus approximans	2	1.60%
Tawny Frogmouth	Podargus strigoides	1	0.80%
Topknot Pigeon	Lopholaimus antarcticus	8	6.40%
Tree Martin	Petrochelidon nigricans	1	0.80%
Variiegated Fairy-wren	Malurus lamberti	24	19.20%
Wedge-tailed Eagle	Aquila audax	8	6.40%
Weebill	Smicrornis brevirostris	1	0.80%
Welcome Swallow	Hirundo neoxena	9	7.20%
Whistling Kite	Haliastur sphenurus	4	3.20%
White-bellied Sea-Eagle	Haliaeetus leucogaster	21	16.80%
White-browed Scrubwren	Sericornis frontalis	62	49.60%
White-cheeked Honeyeater	Phylidonyris niger	19	15.20%
White-faced Heron	Egretta novaehollandiae	1	0.80%
White-naped Honeyeater	Melithreptus lunatus	15	12.00%
White-throated Gerygone	Gerygone olivacea	1	0.80%
White-throated Needle-tail	Hirundapus caudacutus	1	0.80%
White-throated Treecreeper	Cormobates leucophaea	51	40.80%
Willie Wagtail	Rhipidura leucophrys	2	1.60%
Wonga Pigeon	Leucosarcia melanoleuca	4	3.20%
Yellow Thornbill	Acanthiza nana	3	2.40%
Yellow-faced Honeyeater	Caligavis chrysops	75	60.00%
Yellow-tailed Black-Cockatoo	Zanda funereus	18	14.40%

Yellow-throated Scrubwren	<i>Sericornis citreogularis</i>	1	0.80%
Yellow-tufted Honeyeater	<i>Lichenostomus melanops</i>	2	1.60%