

Historical Species List for Narrawallee Inlet from Birdata

Common Name	Scientific Name	Count	Reporting Rate
Australasian Darter	<i>Anhinga novaehollandiae</i>	5	3.73%
Australasian Figbird	<i>Sphecotheres vieilloti</i>	3	2.24%
Australasian Gannet	<i>Morus serrator</i>	6	4.48%
Australasian Grebe	<i>Tachybaptus novaehollandiae</i>	1	0.75%
Australian King-Parrot	<i>Alisterus scapularis</i>	34	25.37%
Australian Magpie	<i>Gymnorhina tibicen</i>	25	18.66%
Australian Pelican	<i>Pelecanus conspicillatus</i>	34	25.37%
Australian Pied Oystercatcher	<i>Haematopus longirostris</i>	46	34.33%
Australian Raven	<i>Corvus coronoides</i>	19	14.18%
Australian White Ibis	<i>Threskiornis moluccus</i>	6	4.48%
Azure Kingfisher	<i>Ceyx azureus</i>	15	11.19%
Bar-tailed Godwit	<i>Limosa lapponica</i>	3	2.24%
Black Swan	<i>Cygnus atratus</i>	3	2.24%
Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>	22	16.42%
Black-faced Monarch	<i>Monarcha melanopsis</i>	12	8.96%
Brown Cuckoo-Dove	<i>Macropygia phasianella</i>	16	11.94%
Brown Falcon	<i>Falco berigora</i>	1	0.75%
Brown Gerygone	<i>Gerygone mouki</i>	39	29.10%
Brown Goshawk	<i>Accipiter fasciatus</i>	2	1.49%
Brown Thornbill	<i>Acanthiza pusilla</i>	63	47.01%
Brush Cuckoo	<i>Cacomantis variolosus</i>	1	0.75%
Caspian Tern	<i>Hydroprogne caspia</i>	5	3.73%
Cattle Egret	<i>Bubulcus ibis</i>	3	2.24%
Channel-billed Cuckoo	<i>Scythrops novaehollandiae</i>	1	0.75%
Chestnut Teal	<i>Anas castanea</i>	2	1.49%
Cicadabird	<i>Edolisoma tenuirostris</i>	1	0.75%
Common Blackbird	<i>Turdus merula</i>	16	11.94%
Common Bronzewing	<i>Phaps chalcoptera</i>	2	1.49%
Common Myna	<i>Acridotheres tristis</i>	3	2.24%
Crested Pigeon	<i>Ocyphaps lophotes</i>	9	6.72%
Crested Shrike-tit	<i>Falcunculus frontatus</i>	4	2.99%
Crested Tern	<i>Thalasseus bergii</i>	20	14.93%
Crimson Rosella	<i>Platycercus elegans</i>	90	67.16%
Crow & Raven spp		1	0.75%
Dollarbird	<i>Eurystomus orientalis</i>	35	26.12%
Domestic Duck		2	1.49%
Dusky Woodswallow	<i>Artamus cyanopterus</i>	2	1.49%
Eastern Curlew	<i>Numenius madagascariensis</i>	9	6.72%
Eastern Koel	<i>Eudynamys orientalis</i>	4	2.99%
Eastern Rosella	<i>Platycercus eximius</i>	2	1.49%
Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>	58	43.28%
Eastern Whipbird	<i>Psophodes olivaceus</i>	55	41.04%
Eastern Yellow Robin	<i>Eopsaltria australis</i>	61	45.52%
Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>	10	7.46%
Galah	<i>Eolophus roseicapilla</i>	26	19.40%
Gang-gang Cockatoo	<i>Callocephalon fimbriatum</i>	4	2.99%
Glossy Black-Cockatoo	<i>Calyptorhynchus lathami</i>	1	0.75%
Golden Whistler	<i>Pachycephala pectoralis</i>	36	26.87%
Great Cormorant	<i>Phalacrocorax carbo</i>	34	25.37%
Great Egret	<i>Ardea alba</i>	18	13.43%
Grey Butcherbird	<i>Cracticus torquatus</i>	21	15.67%
Grey Fantail	<i>Rhipidura fuliginosa</i>	63	47.01%
Grey Goshawk	<i>Accipiter novaehollandiae</i>	2	1.49%
Grey Shrike-thrush	<i>Colluricincla harmonica</i>	34	25.37%
Hooded Plover	<i>Thinornis cucullatus</i>	3	2.24%
Intermediate Egret	<i>Ardea intermedia</i>	2	1.49%
Jacky Winter	<i>Microeca fascinans</i>	1	0.75%

Large-billed Scrubwren	<i>Sericornis magnirostra</i>	5	3.73%
Laughing Kookaburra	<i>Dacelo novaeguineae</i>	76	56.72%
Leaden Flycatcher	<i>Myiagra rubecula</i>	12	8.96%
Lesser Frigatebird	<i>Fregata ariel</i>	1	0.75%
Lewin's Honeyeater	<i>Meliphaga lewinii</i>	65	48.51%
Lewin's Rail	<i>Lewinia pectoralis</i>	1	0.75%
Little Black Cormorant	<i>Phalacrocorax sulcirostris</i>	45	33.58%
Little Corella	<i>Cacatua sanguinea</i>	1	0.75%
Little Eagle	<i>Hieraaetus morphnoides</i>	2	1.49%
Little Egret	<i>Egretta garzetta</i>	8	5.97%
Little Lorikeet	<i>Glossopsitta pusilla</i>	2	1.49%
Little Penguin	<i>Eudyptula minor</i>	1	0.75%
Little Pied Cormorant	<i>Microcarbo melanoleucos</i>	54	40.30%
Little Tern	<i>Sternula albifrons</i>	1	0.75%
Little Wattlebird	<i>Anthochaera chrysoptera</i>	62	46.27%
Magpie-lark	<i>Grallina cyanoleuca</i>	40	29.85%
Masked Lapwing	<i>Vanellus miles</i>	30	22.39%
Mistletoebird	<i>Dicaeum hirundinaceum</i>	13	9.70%
Musk Lorikeet	<i>Glossopsitta concinna</i>	8	5.97%
Nankeen Kestrel	<i>Falco cenchroides</i>	1	0.75%
New Holland Honeyeater	<i>Phylidonyris novaehollandiae</i>	51	38.06%
Noisy Friarbird	<i>Philemon corniculatus</i>	18	13.43%
Noisy Miner	<i>Manorina melanocephala</i>	1	0.75%
Olive-backed Oriole	<i>Oriolus sagittatus</i>	8	5.97%
Osprey	<i>Pandion haliaetus</i>	1	0.75%
Peaceful Dove	<i>Geopelia placida</i>	1	0.75%
Peregrine Falcon	<i>Falco peregrinus</i>	1	0.75%
Pied Cormorant	<i>Phalacrocorax varius</i>	6	4.48%
Pied Currawong	<i>Strepera graculina</i>	26	19.40%
Rainbow Bee-eater	<i>Merops ornatus</i>	1	0.75%
Rainbow Lorikeet	<i>Trichoglossus moluccanus</i>	98	73.13%
Red Wattlebird	<i>Anthochaera carunculata</i>	71	52.99%
Red-browed Finch	<i>Neochmia temporalis</i>	33	24.63%
Red-browed Treecreeper	<i>Climacteris erythropis</i>	3	2.24%
Rose Robin	<i>Petroica rosea</i>	6	4.48%
Rufous Fantail	<i>Rhipidura rufifrons</i>	11	8.21%
Rufous Whistler	<i>Pachycephala rufiventris</i>	22	16.42%
Sacred Kingfisher	<i>Todiramphus sanctus</i>	24	17.91%
Satin Bowerbird	<i>Ptilonorhynchus violaceus</i>	30	22.39%
Satin Flycatcher	<i>Myiagra cyanoleuca</i>	1	0.75%
Scaly-breasted Lorikeet	<i>Trichoglossus chlorolepidotus</i>	1	0.75%
Scarlet Honeyeater	<i>Myzomela sanguinolenta</i>	15	11.19%
Shining Bronze-Cuckoo	<i>Chalcites lucidus</i>	6	4.48%
Short-tailed Shearwater	<i>Ardena tenuirostris</i>	1	0.75%
Silver Gull	<i>Chroicocephalus novaehollandiae</i>	48	35.82%
Silvereye	<i>Zosterops lateralis</i>	23	17.16%
Sooty Oystercatcher	<i>Haematopus fuliginosus</i>	11	8.21%
Sooty Shearwater	<i>Ardena grisea</i>	1	0.75%
Southern Boobook	<i>Ninox boobook</i>	1	0.75%
Southern Emu-wren	<i>Stipiturus malachurus</i>	6	4.48%
Spangled Drongo	<i>Dicrurus bracteatus</i>	1	0.75%
Spotted Dove	<i>Streptopelia chinensis</i>	21	15.67%
Spotted Pardalote	<i>Pardalotus punctatus</i>	25	18.66%
Square-tailed Kite	<i>Lophoictinia isura</i>	1	0.75%
Striated Heron	<i>Butorides striata</i>	12	8.96%
Striated Pardalote	<i>Pardalotus striatus</i>	1	0.75%
Striated Thornbill	<i>Acanthiza lineata</i>	27	20.15%
Sulphur-crested Cockatoo	<i>Cacatua galerita</i>	2	1.49%
Superb Fairy-wren	<i>Malurus cyaneus</i>	42	31.34%

Tawny Frogmouth	Podargus strigoides	2	1.49%
Topknot Pigeon	Lopholaimus antarcticus	1	0.75%
Varied Sittella	Daphoenositta chrysoptera	3	2.24%
Variegated Fairy-wren	Malurus lamberti	27	20.15%
Weebill	Smicrornis brevirostris	1	0.75%
Welcome Swallow	Hirundo neoxena	31	23.13%
Whimbrel	Numenius phaeopus	15	11.19%
Whistling Kite	Haliastur sphenurus	8	5.97%
White-bellied Cuckoo-shrike	Coracina papuensis	1	0.75%
White-bellied Sea-Eagle	Haliaeetus leucogaster	50	37.31%
White-breasted Woodswallow	Artamus leucorhynchus	7	5.22%
White-browed Scrubwren	Sericornis frontalis	28	20.90%
White-browed Woodswallow	Artamus superciliosus	2	1.49%
White-cheeked Honeyeater	Phylidonyris niger	1	0.75%
White-eared Honeyeater	Nesoptilotis leucotis	2	1.49%
White-faced Heron	Egretta novaehollandiae	65	48.51%
White-headed Pigeon	Columba leucomela	13	9.70%
White-naped Honeyeater	Melithreptus lunatus	8	5.97%
White-necked Heron	Ardea pacifica	2	1.49%
White-throated Needletail	Hirundapus caudacutus	1	0.75%
White-throated Treecreeper	Cormobates leucophaea	46	34.33%
Willie Wagtail	Rhipidura leucophrys	27	20.15%
Wonga Pigeon	Leucosarcia melanoleuca	1	0.75%
Yellow Thornbill	Acanthiza nana	25	18.66%
Yellow-faced Honeyeater	Caligavis chrysops	75	55.97%
Yellow-rumped Thornbill	Acanthiza chrysorrhoa	1	0.75%
Yellow-tailed Black-Cockatoo	Zanda funereus	18	13.43%