

Historical Species List for Yatte Yattah from Birdata

Common Name	Scientific Name	Count	Reporting Rate
Australasian Figbird	<i>Sphecotheres vieillotii</i>	1	1.45%
Australian Hobby	<i>Falco longipennis</i>	1	1.45%
Australian King-Parrot	<i>Alisterus scapularis</i>	10	14.49%
Australian Magpie	<i>Gymnorhina tibicen</i>	39	56.52%
Australian Raven	<i>Corvus coronoides</i>	27	39.13%
Australian Wood Duck	<i>Chenonetta jubata</i>	8	11.59%
Barn Owl	<i>Tyto alba</i>	1	1.45%
Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>	16	23.19%
Black-faced Monarch	<i>Monarcha melanopsis</i>	20	28.99%
Black-shouldered Kite	<i>Elanus axillaris</i>	2	2.90%
Brown Cuckoo-Dove	<i>Macropygia phasianella</i>	17	24.64%
Brown Falcon	<i>Falco berigora</i>	3	4.35%
Brown Gerygone	<i>Gerygone mouki</i>	27	39.13%
Brown Quail	<i>Synoicus ypsilophora</i>	3	4.35%
Brown Thornbill	<i>Acanthiza pusilla</i>	23	33.33%
Brown-headed Honeyeater	<i>Melithreptus brevirostris</i>	5	7.25%
Brush Cuckoo	<i>Cacomantis variolosus</i>	2	2.90%
Cattle Egret	<i>Bubulcus ibis</i>	4	5.80%
Channel-billed Cuckoo	<i>Scythrops novaehollandiae</i>	4	5.80%
Chestnut Teal	<i>Anas castanea</i>	3	4.35%
Cicadabird	<i>Edolisoma tenuirostris</i>	1	1.45%
Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>	2	2.90%
Common Blackbird	<i>Turdus merula</i>	1	1.45%
Common Bronzewing	<i>Phaps chalcoptera</i>	1	1.45%
Common Myna	<i>Acridotheres tristis</i>	3	4.35%
Common Starling	<i>Sturnus vulgaris</i>	20	28.99%
Crested Pigeon	<i>Ocyphaps lophotes</i>	2	2.90%
Crested Shrike-tit	<i>Falcunculus frontatus</i>	7	10.14%
Crimson Rosella	<i>Platycercus elegans</i>	38	55.07%
Dollarbird	<i>Eurystomus orientalis</i>	15	21.74%
Dusky Woodswallow	<i>Artamus cyanopterus</i>	2	2.90%
Eastern Koel	<i>Eudynamis orientalis</i>	1	1.45%
Eastern Rosella	<i>Platycercus eximius</i>	31	44.93%
Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>	20	28.99%
Eastern Whipbird	<i>Psophodes olivaceus</i>	24	34.78%
Eastern Yellow Robin	<i>Eopsaltria australis</i>	40	57.97%
Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>	13	18.84%
Galah	<i>Eolophus roseicapilla</i>	21	30.43%
Golden Whistler	<i>Pachycephala pectoralis</i>	33	47.83%
Green Catbird	<i>Ailuroedus crassirostris</i>	4	5.80%
Grey Butcherbird	<i>Cracticus torquatus</i>	34	49.28%
Grey Fantail	<i>Rhipidura fuliginosa</i>	53	76.81%
Grey Goshawk	<i>Accipiter novaehollandiae</i>	1	1.45%
Grey Shrike-thrush	<i>Colluricincla harmonica</i>	34	49.28%
Jacky Winter	<i>Microeca fascinans</i>	31	44.93%
Laughing Kookaburra	<i>Dacelo novaeguineae</i>	35	50.72%
Leaden Flycatcher	<i>Myiagra rubecula</i>	10	14.49%
Lewin's Honeyeater	<i>Meliphaga lewinii</i>	48	69.57%
Little Corella	<i>Cacatua sanguinea</i>	8	11.59%
Little Eagle	<i>Hieraaetus morphnoides</i>	1	1.45%
Little Wattlebird	<i>Anthochaera chrysoptera</i>	5	7.25%
Magpie-lark	<i>Grallina cyanoleuca</i>	18	26.09%
Masked Lapwing	<i>Vanellus miles</i>	4	5.80%
Mistletoebird	<i>Dicaeum hirundinaceum</i>	5	7.25%
Nankeen Kestrel	<i>Falco cenchroides</i>	4	5.80%
New Holland Honeyeater	<i>Phylidonyris novaehollandiae</i>	2	2.90%
Noisy Friarbird	<i>Philemon corniculatus</i>	7	10.14%

Olive-backed Oriole	<i>Oriolus sagittatus</i>	9	13.04%
Painted Button-quail	<i>Turnix varius</i>	1	1.45%
Pallid Cuckoo	<i>Heteroscenes pallidus</i>	8	11.59%
Peregrine Falcon	<i>Falco peregrinus</i>	2	2.90%
Pied Currawong	<i>Strepera graculina</i>	19	27.54%
Rainbow Lorikeet	<i>Trichoglossus moluccanus</i>	33	47.83%
Red Wattlebird	<i>Anthochaera carunculata</i>	28	40.58%
Red-browed Finch	<i>Neochmia temporalis</i>	19	27.54%
Rose Robin	<i>Petroica rosea</i>	2	2.90%
Rufous Fantail	<i>Rhipidura rufifrons</i>	9	13.04%
Rufous Whistler	<i>Pachycephala rufiventris</i>	25	36.23%
Sacred Kingfisher	<i>Todiramphus sanctus</i>	11	15.94%
Satin Bowerbird	<i>Ptilonorhynchus violaceus</i>	15	21.74%
Satin Flycatcher	<i>Myiagra cyanoleuca</i>	1	1.45%
Scarlet Honeyeater	<i>Myzomela sanguinolenta</i>	2	2.90%
Shining Bronze-Cuckoo	<i>Chalcites lucidus</i>	1	1.45%
Silveryeye	<i>Zosterops lateralis</i>	14	20.29%
Southern Boobook	<i>Ninox boobook</i>	1	1.45%
Spectacled Monarch	<i>Symposiachrus trivirgatus</i>	1	1.45%
Spotted Dove	<i>Streptopelia chinensis</i>	2	2.90%
Spotted Pardalote	<i>Pardalotus punctatus</i>	16	23.19%
Straw-necked Ibis	<i>Threskiornis spinicollis</i>	2	2.90%
Striated Pardalote	<i>Pardalotus striatus</i>	2	2.90%
Striated Thornbill	<i>Acanthiza lineata</i>	22	31.88%
Sulphur-crested Cockatoo	<i>Cacatua galerita</i>	7	10.14%
Superb Fairy-wren	<i>Malurus cyaneus</i>	28	40.58%
Superb Lyrebird	<i>Menura novaehollandiae</i>	7	10.14%
Swamp Harrier	<i>Circus approximans</i>	1	1.45%
Topknot Pigeon	<i>Lopholaimus antarcticus</i>	23	33.33%
Varied Sittella	<i>Daphoenositta chrysoptera</i>	10	14.49%
Variegated Fairy-wren	<i>Malurus lamberti</i>	1	1.45%
Wedge-tailed Eagle	<i>Aquila audax</i>	4	5.80%
Welcome Swallow	<i>Hirundo neoxena</i>	7	10.14%
Whistling Kite	<i>Haliastur sphenurus</i>	1	1.45%
White-bellied Cuckoo-shrike	<i>Coracina papuensis</i>	3	4.35%
White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>	3	4.35%
White-browed Scrubwren	<i>Sericornis frontalis</i>	5	7.25%
White-eared Honeyeater	<i>Nesoptilotis leucotis</i>	1	1.45%
White-faced Heron	<i>Egretta novaehollandiae</i>	2	2.90%
White-headed Pigeon	<i>Columba leucomela</i>	3	4.35%
White-naped Honeyeater	<i>Melithreptus lunatus</i>	2	2.90%
White-necked Heron	<i>Ardea pacifica</i>	3	4.35%
White-throated Gerygone	<i>Gerygone olivacea</i>	6	8.70%
White-throated Needletail	<i>Hirundapus caudacutus</i>	1	1.45%
White-throated Treecreeper	<i>Cormobates leucophaea</i>	11	15.94%
White-winged Chough	<i>Corcorax melanorhamphos</i>	1	1.45%
Willie Wagtail	<i>Rhipidura leucophrys</i>	26	37.68%
Wonga Pigeon	<i>Leucosarcia melanoleuca</i>	18	26.09%
Yellow Thornbill	<i>Acanthiza nana</i>	3	4.35%
Yellow-faced Honeyeater	<i>Caligavis chrysops</i>	33	47.83%
Yellow-rumped Thornbill	<i>Acanthiza chrysorrhoa</i>	1	1.45%
Yellow-tailed Black-Cockatoo	<i>Zanda funereus</i>	6	8.70%